

Establishment of Intestinal Identity and Epithelial-Mesenchymal Signaling by *Cdx2*

Nan Gao,¹ Peter White,¹ and Klaus H. Kaestner^{1,*}

¹Department of Genetics, and Institute for Diabetes, Obesity and Metabolism, University of Pennsylvania, Philadelphia, PA 19104, USA

*Correspondence: kaestner@mail.med.upenn.edu

DOI 10.1016/j.devcel.2009.02.010

SUMMARY

We demonstrate that conditional ablation of the homeobox transcription factor *Cdx2* from early endoderm results in the replacement of the posterior intestinal epithelium with keratinocytes, a dramatic cell fate conversion caused by ectopic activation of the foregut/esophageal differentiation program. This anterior homeotic transformation of the intestine was first apparent in the early embryonic *Cdx2*-deficient gut by a caudal extension of the expression domains of several key foregut endoderm regulators. While the intestinal transcriptome was severely affected, *Cdx2* deficiency only transiently modified selected posterior *Hox* genes and the primary enteric *Hox* code was maintained. Further, we demonstrate that *Cdx2*-directed intestinal cell fate adoption plays an important role in the establishment of normal epithelial-mesenchymal interactions, as multiple signaling pathways involved in this process were severely affected. We conclude that *Cdx2* controls important aspects of intestinal identity and development, and that this function is largely independent of the enteric *Hox* code.

INTRODUCTION

The mouse endoderm transforms from a two-dimensional epithelial sheet into the primitive gut tube at embryonic day 8.5–9.0 (E8.5–E9.0). Subsequent morphological differentiation converts the pseudostratified endoderm layer into a tall columnar epithelium which lines the respiratory and gastrointestinal tracts (Wells and Melton, 1999). The primitive gut appears homogeneous from end to end, with distinct anterior-posterior (AP) regions adopting different fates in subsequent organogenesis. In the gastrointestinal tract, the foregut gives rise to the epithelia of esophagus, stomach, and duodenum, while midgut and hindgut become the small intestine, and the cecum and colon, respectively. Crosstalk between gut mesoderm and endoderm progressively commits the primary endoderm to specific fates (Grapin-Botton and Melton, 2000). Mutations in a number of *Hox* genes result in malformations in certain gut regions, but do not cause wholesale AP transformation of the gut (Aubin et al., 1997; Boulet and Capecchi, 1996; Manley and Capecchi, 1995; Warot et al., 1997; Zacchetti et al., 2007), even though these *Hox* factors play important roles in patterning

the mesoderm and neural tube (Deschamps et al., 1999; Krumlauf, 1994; McGinnis and Krumlauf, 1992).

AP asymmetry of the gut endoderm is evident with the onset of *Cdx2* expression in the hindgut at its inception (Beck et al., 1995). *Cdx2* is the mouse homolog of *AmphiCdx* in *Amphioxus* and *caudal* in *Drosophila*. It resides in the “*ParaHox*” gene cluster believed to have evolved from a “*ProtoHox*” cluster that also gave rise to the definitive *Hox* gene clusters (Brooke et al., 1998). In *Drosophila*, *caudal* specifies posterior body segments (Macdonald and Struhl, 1986; Mlodzik et al., 1985; Moreno and Morata, 1999). Morpholino knockdown and overexpression studies in zebrafish indicated essential roles of *caudal* orthologs in neural tube and intestinal development (Cheng et al., 2008; Flores et al., 2008; Shimizu et al., 2006; Skromne et al., 2007). Three mouse homologs, *Cdx1*, *Cdx2*, and *Cdx4* (Duprey et al., 1988; Gamer and Wright, 1993; James and Kazenwadel, 1991), participate in the patterning of the vertebral column (Chawengsaksophak et al., 1997; Subramanian et al., 1995; van Nes et al., 2006) and in embryonic hematopoiesis (Wang et al., 2008); however, their role in endoderm development is less clear. Homozygous *Cdx1* or *Cdx4* mutants do not display gut defects (Subramanian et al., 1995; van Nes et al., 2006), while *Cdx2* null mutants die in utero before the onset of endoderm development (Chawengsaksophak et al., 1997; Tamai et al., 1999).

In the mouse embryo, *Cdx2* is expressed in nuclei of cells derived from the late-dividing blastomere, a precursor of trophoblast (Deb et al., 2006). From E8.5 onward, *Cdx2* is activated in the embryo proper, predominantly the posterior gut (Beck et al., 1995). *Cdx2* expression subsequently becomes restricted to the intestinal epithelium, with a sharp anterior boundary marking the transition from stomach to duodenum (James et al., 1994; Silberg et al., 2000). Genetic analysis of *Cdx2* function in mammalian intestinal development has been limited to *Cdx2* heterozygous mice that form multiple colonic polyps (Chawengsaksophak et al., 1997). These polyps contain areas of squamous metaplasia in which the expression of the remaining wild-type *Cdx2* allele is extinguished through an unknown mechanism (Beck et al., 1999). However, it is still unclear as to (1) what transcriptional programs were altered in these epimorphic lesions; (2) how comprehensive the impact of *Cdx2* disruption is for cell fate determination; (3) through which mechanisms loss of *Cdx2* induces squamous metaplasia; (4) through which mechanisms *Cdx2* promotes intestinal differentiation; and (5) what role *Cdx2* plays in intestinal epithelial-mesenchymal interactions.

We have previously shown that ectopic expression of *Cdx2* in the gastric epithelium induces intestinal metaplasia (Silberg et al., 2002), an example of a posterior homeotic transformation.




Figure 1. Conditional Ablation of *Cdx2* in Mouse Endoderm Leads to Abnormal Intestinal Growth and Colon Dysgenesis

(A and B) Whole-mount β-galactosidase staining of E9.5 *Cdx2*^{loxP/+}, *Foxa3Cre*, *R26R* and *Cdx2*^{loxP/loxP}, *Foxa3Cre*, *R26R* embryos. (C and D) Immunohistochemistry for *Cdx2* in E9.5 control and mutant embryos. (E and F) A pan endoderm marker, *Foxa1*, continues to be expressed in the *Cdx2*-deficient endoderm. (G) The distal intestine of the E12.5 mutant gut forms a blind-ended sac (arrow). (H) At E14.5, abnormal intestinal growth and terminal blockage are evident in the mutant posterior gut, with no colon formation. Arrow points to malformed cecum. (I–N) Confocal immunostaining for β-catenin (green) and *Cdx2* (red) was performed on E14.5 anterior (I and J), medial (K and L), and posterior (M and N) intestinal sections. Nuclei were counterstained by DAPI in blue. Scale bars: 50 μm.

resistance gene was removed by crossing to *Flp1* deleter mice (Rodriguez et al., 2000). *Cdx2*^{loxP/+} mice were then intercrossed, resulting in *Cdx2*^{loxP/loxP} mice that were viable and fertile (Figure S1C), confirming that the *Cdx2*^{loxP} allele is functionally wild-type. Subsequent Cre-mediated gene ablation results in a null allele that lacks the homeobox domain. To ablate *Cdx2* conditionally in the developing gut, we bred *Cdx2*^{loxP/+} mice to *Foxa3Cre* mice (Lee et al., 2005), which delete loxP-flanked targets in early endoderm. Using the *Rosa26R* reporter line, we verified Cre activity in the primitive

gut of E9.5 embryos, prior to the onset of gross morphological defects (Figures 1A and 1B). Efficient deletion of *Cdx2* from mutant (*Cdx2*^{loxP/loxP}, *Foxa3Cre*⁺) gut epithelia was evident with immunohistochemistry using an anti-*Cdx2* antibody (Figures 1C and 1D). The expression of *Foxa1*, a pan-endoderm marker, was unaffected (Figures 1E and 1F). Examination of mutant embryos at mid and late gestation revealed equal efficiency of *Cdx2* ablation throughout the intestinal domain (Figures 1I–1N and Figure S2).

Here, we demonstrate that *Cdx2* is essential for the initial expression and/or subsequent maintenance of a group of prointestinal transcription factors, including *Cdx1*, *Isx*, *HNF1α*, and *HNF4α*, which together activate the intestinal transcriptome. The expression of *Cdx2* in the posterior gut epithelium antagonizes the foregut differentiation program, which becomes ectopically activated upon *Cdx2* disruption, resulting in dramatic cell fate conversion. We further demonstrate that intestinal cell fate establishment by *Cdx2* plays a critical role in instructing normal epithelial and mesenchymal interactions, in particular with respect to the integrity of Wnt and Hedgehog signaling.

RESULTS

Conditional Ablation of *Cdx2* from the Developing Endoderm

Cdx2 null mice die before gastrulation (Chawengsaksophak et al., 1997; Tamai et al., 1999). Therefore, we derived a conditional *Cdx2* allele to study its role in the gut endoderm (see Figure S1A available online). Correctly targeted embryonic stem cell clones were identified by Southern blot analysis (Figure S1B). After germline transmission of the targeted allele, the FRT-flanked neomycin

gut of E9.5 embryos, prior to the onset of gross morphological defects (Figures 1A and 1B). Efficient deletion of *Cdx2* from mutant (*Cdx2*^{loxP/loxP}, *Foxa3Cre*⁺) gut epithelia was evident with immunohistochemistry using an anti-*Cdx2* antibody (Figures 1C and 1D). The expression of *Foxa1*, a pan-endoderm marker, was unaffected (Figures 1E and 1F). Examination of mutant embryos at mid and late gestation revealed equal efficiency of *Cdx2* ablation throughout the intestinal domain (Figures 1I–1N and Figure S2).

Intestinal Growth Is Severely Affected in *Cdx2* Mutant Mice

Although the mutant pups were born alive, they did not survive beyond postnatal day one (P1). We examined the gastrointestinal tract of mutant embryos at various developmental stages. The gross abnormalities of the mutant posterior gut region first became evident at E12.5 (Figure 1G). In contrast to the control intestinal tract that ends with colon and rectum, the mutant intestine developed an abnormal distal structure that terminates in a blind-ended sac (Figure 1G, arrow). Progressive defects in elongation of the mutant intestine began to appear at E14.5 (Figure S3A), and the mutant gastrointestinal tract developed a malformed cecum at the distal end, with no colon (Figure 1H,




Figure 2. Initial Intestinal Differentiation Is Blocked in the Cdx2-Deficient Gut


(A and B) Severe intestinal shortening is evident in the E16.5 mutant duodenum. The colon fails to form in Cdx2 mutant embryos. Arrows point to cecum. d, duodenum; p, pancreas; s, stomach; sp, spleen.
(C–D') E18.5 mutant intestine is dilated due to terminal blockage. duo, duodenum; jej, jejunum; ile, ileum.
(E and F) Alkaline phosphatase staining (red) of E16.5 control and mutant jejunum sections.
(G and H) Confocal immunostaining for p63 (red) and DBA lectin (green) of E15.5 control and mutant jejunum sections. Nuclei were counterstained by DAPI in blue.
(I and J) Alkaline phosphatase staining (red) of E18.5 control and Cdx2 mutant jejunum sections demonstrates absence of goblet cells in mutant epithelium.
(K and L) Alcian blue staining (blue) of E18.5 control and Cdx2 mutant jejunum sections demonstrates absence of goblet cells in mutant epithelium.
(M and N) Confocal immunostaining of Cdx2 (green), lectin DBA (red) and chromogranin A (blue) demonstrates absence of lectin-positive intestinal mucosa as well as enteroendocrine cells in Cdx2 mutants. Arrowheads in (M) point to the rare enteroendocrine cells in the control intestine.
Scale bars: 75 μ m in (E), (F), and (I)–(L); 50 μ m in (G), (H), (I'), (K'), (M), and (N).

arrow). Cross-sections of the E14.5 mutant distal intestine revealed a dilated gut lumen (Figures 1M and 1N). All mutant animals examined from E13 to P0 ($n = 56$) demonstrated an absence of the colon, a phenotype reminiscent of the most severe cases of colonic atresia in humans (Etensel et al., 2005; Lau and Caty, 2006). The mutant duodenum was progressively distended and became translucent, likely due to fluid retention caused by distal obstruction (Figures 2A and 2B). By E18.5, the duodenum was further dilated to a 5- to 7-fold increase in diameter compared to the control tissue (Figures 2C, 2D, and 2D'). Thus, Cdx2 deficiency prevents colon formation and leads to complete intestinal obstruction.

While the mutant proximal and medial intestinal epithelia appeared less organized than the control epithelia, the overall histology at E14.5 differed only subtly, as both mutant and control gut epithelia appeared pseudostratified (Figures 1I–1N). However, defects in differentiation became more apparent later in development. Since the mutant animals die at P1, before the development of Paneth cells, we examined the differentiation of enterocytes (Figures 2E, 2F, 2I, and 2J), goblet cells (Figures 2G, 2H, and 2K–2N), and enteroendocrine cells (Figures 2M

and 2N) at different stages using specific markers and found terminal differentiation severely impaired. Instead, the mutant posterior intestinal epithelium expressed a basal epithelial cell marker p63 from E15.5 onward (Figure 2H).

Villus hypoplasia was detected from E16.5 throughout the mutant intestinal domain as compared to controls (Figures 3A and 3B). Position-matched longitudinal histological sections of E18.5 control and mutant intestines revealed dramatic reductions of intestinal villi (Figures 3C–3J), with severity increasing from anterior to posterior (Figure S3B). The mutant duodenum contained villus-like epithelial foldings (Figure 3D) that were significantly stunted and broadened (Figure 3F and Figure S3C), while the cuboidal epithelia of mutant jejunum and ileum were completely replaced with a flattened epithelium (Figures 3H and 3J), and the mutant ileum and cecum lacked villi entirely (Figure 3J and Figure S2F for cecum). Intestinal epithelia containing mosaic Cdx2 deletion were observed in a few mutant embryos. Segments of Cdx2-positive epithelium were contiguous to Cdx2-deficient regions (Figure 3K). Interestingly, cells that retained Cdx2 expression were capable of forming villi and elaborate goblet cells normally (Figure 3L), while adjacent



Cdx2-deficient cells failed to do so (Figure 3M). Thus, *Cdx2* is required for initiation of intestinal differentiation and morphogenesis in a cell-autonomous fashion.

Excess Proliferation and Keratinocyte Character in the Mutant Intestine

Ki67 staining, which marks transit-amplifying cells, revealed an expanded proliferative compartment in the mutant duodenal epithelium (Figures 3N and 3O). The proliferative index of the mutant duodenal epithelial cells, assayed by BrdU incorporation, was significantly increased in the mutant epithelium following either 1 or 24 hr labeling (Figures S4A–S4C). Interestingly, even after a short labeling period (40–60 min), more than 20% of BrdU⁺ mutant cells were located at or above position 9 relative to the bottom of the nascent crypts (Figures 3Q and 3R and Figure S4D), while BrdU⁺ control cells were restricted to the inter-villus space even after 24 hr BrdU labeling (Figures 3P and

Figure 3. *Cdx2* Deficiency Affects Villus Morphogenesis and Proliferation Pattern

(A and B) Immunohistochemistry for *Cdx2* on E16.5 control and mutant ileal sections.

(C–J) H&E staining of position-matched E18.5 longitudinal sections of control and mutant duodenum, jejunum, and ileum. The mutant duodenal epithelial folding is stunted (F).

(K–M) Mosaic *Cdx2* deletion in E16.5 mutant ileum is indicated by *Cdx2* immunostaining (brown). *Cdx2*⁺ epithelia are contiguous to regions of *Cdx2*[−] epithelia (K). Differentiated goblet cells (Alcian blue) were observed only in the *Cdx2*⁺ epithelial cells (arrowhead in [L] and [M]), but not in the *Cdx2*[−] epithelium.

(N and O) Immunohistochemistry for Ki67 on E18.5 duodenal sections.

(P and Q) BrdU incorporation followed by immunohistochemistry illustrates that proliferating cells are distributed throughout the stunted villi of *Cdx2*-deficient mice, in contrast to control BrdU⁺ cells that are localized to the intervillus region.

(R) Quantification of BrdU⁺ cells located along the crypt-villus axis after 60 min of *in vivo* labeling.

Scale bars: 75 μ m in (A) and (B); 100 μ m in (C), (D), (G), and (K); 50 μ m in (E), (F), and (L)–(Q).

3R and Figure S4E). BrdU incorporation revealed a continuous proliferative cell layer across the mutant epithelial sheet in the posterior intestine (Figure S5B). We did not, however, detect a significant increase in the apoptotic rate in *Cdx2* mutants by either cleaved caspase-3 or TUNEL staining (Figure S5D). Thus, the failure to specify the colon was not caused by a lack of proliferative capacity or enhanced cell death in *Cdx2* mutant mice. The loss of terminal differentiation discussed above, and our failure to observe gastric glandular epithelial cell types using specific antibodies (data not shown), suggest that the proliferative

pattern of the mutant epithelium resembles that of early embryonic stages prior to intestinal differentiation.

To gain insight into the identity of the *Cdx2* mutant epithelial cells, especially those in the posterior intestinal epithelium, we examined their ultrastructural features using transmission electron microscopy. *Cdx2*-deficient epithelial cells failed to develop the brush border typical of enterocytes (Figures 4A–4D). Examination of the posterior intestine revealed multiple layers of flattened epithelial cells, with the axes of the nuclei parallel to the luminal surface (Figure 4F). Unexpectedly, the mutant posterior intestinal epithelial cells contained abundant tonofilaments (Figure 4G). Tonofilaments are typical of squamous epithelial cells and are frequently seen in the desmosomal junctions of keratinocytes (Figure 4H), which contribute to stratified esophageal epithelia but are extremely rare in the normal intestine (Figure 4E).

Squamous differentiation has been reported in colorectal adenoma (Ouban et al., 2002). To verify whether the *Cdx2*-deficient




Figure 4. The *Cdx2*-Deficient Intestine Lacks Intestine-Specific Ultrastructural Features

(A–D) Electron micrographs of E18.5 control and mutant anterior intestinal epithelium. Mutant cells lack brush border (D).

(E and F) Electron micrographs of E18.5 control and mutant posterior intestinal epithelium. Double-headed arrows indicate the orientations of the mutant nuclei which are parallel to the luminal surface (F), as opposed to the perpendicular orientation in control cells (E). Blue arrowheads point to goblet cells in control tissue (E).

(G and H) Mutant ileal cells contain abundant tonofilaments (yellow arrowheads in [G]) across intercellular junctions where desmosome-like structures (yellow arrowheads in [H]) are assembled. Scale bars: 2 microns in (A), (B), (E), and (F); 500 nm in (C), (D), (G), and (H).

posterior intestine has molecular features of squamous epithelia, we performed immunohistochemistry for keratin 13 and p63, markers of the suprabasal and basal squamous epithelial cells in mouse esophagus, respectively (Figures 5A and 5B). In contrast to the control ileum where neither gene was expressed (Figures 5C and 5D), the *Cdx2* mutant epithelium was positive for both markers (Figures 5E–5G). Neither *keratin 13* nor *p63* was expressed in E10.5 wild-type midgut or hindgut endoderm (Figures S6C and S6D). At E12.5, p63 expression was detected in foregut endoderm cells fated to become forestomach and pharynx (Figures S6E and S6F). Likewise, a marker of the anterior foregut endoderm, Sox2 (Que et al., 2007), was detected in the *Cdx2*-deficient ileum at an expression level equivalent to that of normal esophagus (Figure 5H), confirming that the *Cdx2*-deficient posterior intestine was indeed anteriorized. These data indicate that the expression of squamous markers in the mutant prospective intestine was not due to a developmental delay, but rather due to an ectopically activated foregut differentiation program.

Global Activation of Esophageal Genes in the *Cdx2*-Deficient Posterior Gut

We next performed gene expression profiling using RNA samples extracted from total E18.5 control and mutant ileum as well as normal esophagus. The morphological abnormalities of the mutant ileum precluded separation of the epithelium from mesenchyme. Of the 11,738 significantly changed genes, 268 genes demonstrated fold-changes above 50-fold compared to control ileum. Hierarchical clustering showed that the transcriptome of the mutant ileum resembled that of esophagus far more than that of normal ileum (Figure S7A). The similarity between the mutant ileum and control esophagus is highlighted by a heat map assembled from differentially expressed genes (Figure 5I). Consistent with the morphological transformation, virtually all intestine-specific genes were downregulated in the mutant ileum (Figure 5J and Table S1).

Next, we compared our microarray results with several previous intestine gene profiling studies (Bates et al., 2002; Li et al., 2007; Schroder et al., 2006). Among genes that show significant enrichment in E18.5 intestinal epithelium over mesenchyme (Li et al., 2007), 35.3% were significantly altered in *Cdx2*-deficient intestine. Likewise, 39.8% of genes that show enrichment in intestine over stomach (Bates et al., 2002) were significantly

affected in our model. Furthermore, genes with highly specific expression patterns in the differentiated intestinal epithelium (Schroder et al., 2006) were all significantly downregulated in *Cdx2*-deficient mice.

In contrast, many genes involved in keratinocyte differentiation were significantly upregulated in the *Cdx2* mutant ileum (Figure 5K, Table S2, and Figure S7B). Notably, nine out of the ten most highly upregulated genes in the mutant ileum were enriched in normal esophageal epithelium (Table S2). Most of these genes, such as high molecular weight *keratins*, *keratin 5 and 13*, *small proline-rich protein 3*, *calmodulin-like 3*, *cornifelin*, *plakophilin 1*, and *dermokine*, play important roles in keratinocyte cell envelope formation and desmosome assembly. Thus, gene profiling analysis confirmed the transformation of the *Cdx2*-deficient prospective intestinal endoderm domain to an esophageal cell fate.

Cdx2 Deficiency Alters the AP Distribution of Transcriptional Regulators

To gain a mechanistic insight into the cell fate switch in the *Cdx2*-deficient posterior intestine, we analyzed transcriptional regulators known to be crucial in regulating intestinal differentiation. Along with the striking decrease of *Cdx2* mRNA itself, several intestine-enriched transcription factors, including *HNF1 α* , *HNF4 α* , *Isx*, and *Cdx1*, were dramatically reduced in expression in the mutant intestine (Table S1). Next, we performed quantitative reverse transcriptase PCR (qRT-PCR) analysis on developing gastrointestinal tracts at earlier embryonic time points. A significant reduction in mRNA levels of *HNF1 α* , *HNF4 α* , and *Isx* was already evident in the E12.5 mutant gut (Figures 6B–6D). At E14.5, expression of all these factors as well as *Cdx1* was significantly reduced in both proximal and distal intestine (Figures 6A–6D). These data indicate that the reduced expression of these transcription factors at E18.5 was due not to a secondary effect of abnormal development, but to an impairment of the initial activation of these genes.

The expression of *Math1*, a basic helix-loop-helix transcription factor that plays a role in the differentiation of intestinal secretory cell types (Yang et al., 2001), was significantly reduced in mutant distal intestines from E12.5 onward (Figure 6E). Conversely, qRT-PCR of *Sox2* and *Pax9* demonstrated that these foregut-enriched genes were ectopically activated in the mutant posterior intestine as early as E12.5, at a level equivalent to the stomach




Figure 5. The *Cdx2*-Deficient Posterior Intestine Activates Esophageal Genes

(A and B) The E18.5 control esophageal epithelium expresses both keratin 13 and p63 (green). Sections were counterstained by E-cadherin (red) and DAPI (blue) in (B).

(C–F) The E18.5 control ileal epithelium lacks keratin 13 and p63 staining, while the *Cdx2* mutant ileal epithelium expresses both markers.

(G) Confocal immunostaining for keratin 13 (green) and p63 on *Cdx2* mutant ileum.

(H) Western blot for Sox2, a foregut endoderm transcription factor.

(I) Heat maps, generated for genes with at least a 20-fold change in mutant ileum compared to control ileum. Scales of the heat map are log based. The blue and red brackets indicate significantly down- and upregulated genes listed in (J) and (K).

(J) A partial list of intestinal genes (asterisks) that were extinguished in the mutant ileum.

(K) A partial list of esophageal genes (asterisks) that are significantly activated in the mutant ileum. Scale bars: 50 μ m in (A), (B), and (G); 75 μ m in (C)–(F).

Cdx1. Among multiple Cdx binding sites within the 5' upstream region of *HNF1 α* , those located near the transcription initiation site were most conserved from *Xenopus* to human (Figure S8A). Less conserved Cdx binding sites were identified in the *HNF4 α* and *Cdx1* 5' upstream sequence (Figures S8B and S8C). These Cdx sites are occupied by Cdx2 in vivo, as demonstrated by chromatin immunoprecipitation (ChIP) (Figure 6L).

Cdx2 Deficiency Affects Expression of Selected Enteric Hox Genes

In a number of nonendoderm tissues, Cdx factors exert their developmental effect by regulating Hox transcription factors (Charite et al., 1998; Shimizu et al., 2006;

Subramanian et al., 1995; Wang et al., 2008), which are key players in the primary AP patterning process of the vertebrate embryo (Krumlauf, 1994). Overexpression or inactivation of specific Hox genes has been shown to affect gastrointestinal development (Aubin et al., 1997; Boulet and Capecchi, 1996; Kondo et al., 1996; Pollock et al., 1992; Wolgemuth et al., 1989), while a cluster of *Hoxd* genes controls the formation of the ileocecal sphincter (Zakany and Duboule, 1999). Our microarray data indicated that a number of intestine-enriched Hox genes, including *Hoxa5*, *Hoxb5*, *Hoxb6*, *Hoxa7*, and *Hoxb7*, continue to be expressed in the mutant ileum at levels similar to controls (Figure S9). However, *Hoxc9*, a gene expressed in the posterior midgut and hindgut (Grabin-Botton and Melton, 2000; Roberts, 2000), was decreased 6.2-fold in the mutant ileum (Figure S9B).

Next we examined the AP distribution of representative Hox genes in the early gut, where their expression patterns have been well documented (Choi et al., 2006; Grabin-Botton and Melton, 2000; Pitera et al., 1999; Roberts, 2000). Levels of




Figure 6. Cdx2 Deficiency Replaces Pro-Intestinal Factors with Foregut Regulators

(A–I) Quantitative RT-PCR analysis was performed on E12.5 and E14.5 control and mutant stomach (St.), proximal small intestine (P. Int.), distal small intestine (D. Int.), and cecum (C.), using gene specific primers. * $p < 0.05$; # $p < 0.01$. Error bars show SEM. (J) The top diagram illustrates tissue segments of the E14.5 gut, from anterior (A) to posterior (P) end, used in the qRT-PCR analysis. The diagram at the bottom summarizes the anteriorization event that occurred in the mutant intestinal domain. The foregut differentiation program is shown in red while intestinal differentiation is depicted in blue. (K) Western blots for Cdx1, HNF1α, HNF4α, Wnt10a. (L) ChIP assay for Cdx2 occupancy of the HNF1α, HNF4α, and Cdx1 promoters. No enrichment is detected in ChIP samples derived from Cdx2 mutant intestines. * $p < 0.05$. The error bars indicate standard error of the mean.

restricted to the duodenum even in the absence of Cdx2 (Figures S10E and S10F). Furthermore, the expression of *Barx1*, a stomach-specific mesenchymal transcription factor (Kim et al., 2005), also maintained its expression domain in Cdx2 mutant embryos (Figure S10C). These data further support the notion that the Cdx2-deficient gut retained certain AP values.

Cdx2 Deficiency Affects Intestinal Epithelial-Mesenchymal Interactions

We found dysregulation of Wnt ligand expression in the Cdx2-deficient intestine. In addition to the ectopic activation

Hoxa3, *Hoxb3*, *Hoxb4*, *Hoxc4*, *Hoxd4*, *Hoxb5*, *Hoxc5*, *Hoxa7*, and *Hoxb7* mRNA were not significantly changed in the Cdx2 mutant (Figures 7A–7H and Figure S10A). In contrast, at E12.5, *Hoxc8*, *Hoxb9*, *Hoxc9*, *Hoxa13*, and *Hoxd13* mRNA levels were significantly lower in the mutant posterior intestine (Figures 7I–7L and Figure S10B). At E14.5, however, most of these posterior *Hox* genes had recovered to match the levels of the control intestine (Figures 7I–7L and Figure S10B).

Most *Hox* genes analyzed are expressed in the gut mesenchyme (Li et al., 2007), while some, such as *Hoxa3*, *Hoxb4*, *Hoxc5*, *Hoxb9*, *Hoxc9*, *Hoxa13*, and *Hoxd13*, are also active in the epithelium (Grapin-Botton and Melton, 2000; Roberts, 2000). Our results demonstrate that Cdx2 deficiency in the early gut endoderm transiently modifies the expression of selected posterior *Hox* genes, but had no impact on anterior *Hox* genes. While the posterior Cdx2-deficient gut was anteriorized as early as E12.5 (Figures 6G–6J), the maintained expression of *Hoxc9*, *Hoxa13*, and *Hoxd13* in this domain (Figures 7J–7L) indicates that the mutant gut had retained its primary enteric *Hox* code.

In addition, the Cdx2-deficient gut demonstrated normal AP expression of *Pdx1*, a second “*Parahox*” gene, which remained

of *Wnt10a* as a result of anteriorization of the mutant ileum (Figures 6I and 6K), multiple other *Wnts*, as well as the Wnt target genes *CD44*, *cyclin D1*, *Sox9*, and the *Tcf* factors, were significantly upregulated in the Cdx2-deficient intestine (Figures 8A and 8B; Figures S11A–S11D). In contrast to Wnt, expression of *Ihh* and *Sonic hedgehog* (*Shh*) was significantly reduced in the Cdx2-deficient ileum (Figure 8A), consistent with the decreased expression of *Hedgehog-interacting protein* (*Hhip*), a primary hedgehog target expressed by the intestinal mesenchyme (Li et al., 2007). The severely expanded smooth muscle layer we observed in the mutant duodenum (Figure 3F) and jejunum (Figures 8C–8F) may reflect the decreased hedgehog signaling activity, as inhibition of hedgehog signaling in the intestine causes smooth muscle expansion (Madison et al., 2005).

The expression of *desmin*, a marker of smooth muscle progenitors but not of myofibroblasts (Adegbayega et al., 2002), was increased 6.2-fold in the Cdx2-deficient intestine and was accompanied by a significant decrease of several *myosin* genes (Figure S11E), suggesting an altered myogenic process and terminal differentiation of smooth muscle cells in the Cdx2 mutant gut. The *myosin* gene expression profile in the Cdx2




Figure 7. *Cdx2* Deficiency Affects Selected Posterior Enteric *Hox* Genes

(A–L) Quantitative RT-PCR analysis of E12.5 and E14.5 control and mutant stomach (St.), proximal small intestine (P. Int.), distal small intestine (D. Int.), and cecum (C), using gene specific primers. * $p < 0.05$; # $p < 0.01$. Error bars show SEM.

the posterior of the *Cdx2* mutant gut tube (Figure 8G). None of the previously reported mutant mice had such a dramatic impact on AP patterning of the gut (Aubin et al., 1997; Boulet and Capecchi, 1996; Manley and Capecchi, 1995; Warot et al., 1997; Zacchetti et al., 2007). When a cluster of *Hoxd* genes was deleted, no dramatic disruption of intestinal identity was observed, except that induction of the cecum was affected (Zacchetti et al., 2007). The cecum was correctly induced in *Cdx2* mutant embryos, consistent with the observation that the primary enteric *Hox* code was maintained.

Though *Cdx* factors have been proposed to function via regulation of *Hox* gene expression in several nonendoderm tissues (Charite et al., 1998; Subramanian et al., 1995; Wang et al., 2008), the expression of anteriorly localized intestinal *Hox* genes was independent of *Cdx2*. *Cdx2* deficiency transiently delayed the expression of several posterior intestinal *Hox* genes at early embryonic stages; however, these genes maintained their relative AP position. The regulation of posterior *Hox* genes by *Cdx* factors has been reported in zebrafish hindbrain (Shimizu et al., 2006); however, functional rescue by downstream *Hox* factors remains controversial (Skromne et al., 2007). Our findings indicate that *Cdx2* deficiency does not profoundly influence the primary enteric *Hox* code.

mutant ileum highly resembled that of wild-type esophagus (Figure S11E), illustrating a potent epithelial-to-mesenchymal regulatory role that affected smooth muscle differentiation.

DISCUSSION

Cdx2 and the AP Patterning of the Gut

The *Cdx2*-deficient gut displays severe hindgut abnormalities with a failure of colon development and a complete terminal blockage. Partial or complete colonic atresia has been reported as a human congenital disorder (Etensel et al., 2005). Mutations in *PDX1*, a neighboring “*Parahox*” gene, cause pancreatic agenesis in humans (Stoffers et al., 1997). Our findings support the notion that the *Parahox* genes specify regional identities in the vertebrate gut, and suggest further that mutations in *CDX2* or its targets could contribute to colonic atresia in humans.

The expression domains of multiple important foregut regulators, including *Sox2*, *Pax9* (Grapin-Botton and Melton, 2000), *p63* (Glickman et al., 2001), were dramatically extended toward

Cdx2 Regulates Pro-Intestinal Transcription Factors

Cdx1, whose gut expression pattern resembles that of *Cdx2* (Silberg et al., 2000), has the capability to drive intestinal differentiation in a gain-of-function setting (Mutoh et al., 2004). Redundancy between all three *Cdx* proteins has been reported in a number of nonendoderm tissues (van den Akker et al., 2002; van Nes et al., 2006; Wang et al., 2008). Therefore, it was surprising to see the near-complete homeotic transformation of the *Cdx2*-deficient intestine, as some compensation was anticipated. We established that *Cdx1* activation is directly dependent on *Cdx2*. This transcriptional hierarchy between the two *Cdx* genes reflects their sequential expression pattern in the gut endoderm, where *Cdx2* precedes *Cdx1* by a few days (Hu et al., 1993; Meyer and Gruss, 1993; Silberg et al., 2000). In fact, the expression of *Cdx1* starts only when villus morphogenesis and epithelial




Figure 8. *Cdx2* Deficiency Affects Epithelial-Mesenchymal Signaling

(A) The heat map shows alterations in the expression of *Wnt* and *Hedgehog* genes in the *Cdx2* mutant gut. Positive and negative fold changes are shown in red and green numbers, respectively. $n = 3$ for each tissue type.

(B) Western blots demonstrate upregulation of several *Wnt* targets in the *Cdx2* mutant intestine.

(C–F) Confocal immunostaining for smooth muscle actin (SMA). Sections were counterstained for E-cadherin (green) and DAPI (blue).

(G) Expression of *Cdx2* in midgut and hindgut endoderm directs these domains toward the intestinal cell fate. *Cdx2* promotes intestinal identity via a feedforward mechanism involving activation of pro-intestinal transcription factors, while a foregut/esophageal cell differentiation program is repressed by *Cdx2* in the posterior gut (right panel). Removing *Cdx2* from posterior endoderm (bottom left) leads to replacement of intestinal epithelial identity by ectopic foregut epithelial differentiation. *Cdx2* is critical for the expression of signaling molecules, the epithelial-mesenchymal interaction, and the intestinal proliferation pattern.

notion that *Cdx2* functions upstream of a group of pro-intestinal transcription factors, with which it synergizes to promote intestinal cell fate (Figure 8G).

***Cdx2* Antagonizes the Foregut Differentiation Program**

Our conditional *Cdx2* mutants recapitulate the finding of squamous metaplasia in *Cdx2*^{+/−} mouse colonic lesions, where *Cdx2* expression was diminished (Beck et al., 1999). When *Cdx2* was removed in our model, anteriorization was first evident with the caudal extension of the Sox2 and Pax9 expression domains (Figure 8G).

This was followed by squamous differentiation around E14.5–E15.5, leading to genome-wide activation of esophageal genes. Due to the lack of pro-intestinal regulators, the prospective intestinal epithelial domain was replaced by keratinocytes. These data provide molecular evidence that *Cdx2* normally represses a foregut differentiation program in the posterior gut, explaining the epimorphic changes observed previously (Beck et al., 1999).

We demonstrate that the normal gastrointestinal expression domain of *Wnt10a* is opposite to that of *Cdx2*, mimicking the expression pattern of Sox2. *Cdx2* deficiency led to ectopic activation of *Wnt10a* expression in the caudal intestine, possibly as a consequence of the ectopically differentiated squamous cells. Recent findings suggest that ectodermal dysplasia in humans is associated with *Wnt10a* mutations (Adaimy et al., 2007), while misregulation of *Wnt10a* was found in gastrointestinal cancer (Kirikoshi et al., 2001). We speculate that this gene may be involved in keratinocyte differentiation during upper gastrointestinal development.

maturation begin (Hu et al., 1993). Our data provide further evidence for the evolutionary significance of the “*Parahox*” cluster, where *Cdx2*, but not *Cdx1* or *Cdx4*, is located. Thus, *Cdx1* is controlled by the more ancient *caudal* ortholog *Cdx2* in gut endoderm to facilitate the developmental and anatomical complexity of the organ.

Similar to *Cdx1*, *Isx* is another intestine-specific transcription factor whose expression initiates during epithelial differentiation (Choi et al., 2006), consistent with its dependency on *Cdx2*. In addition, the maintenance of *HNF1α* and *HNF4α* expression in the embryonic intestine is directly controlled by *Cdx2*. Single-gene ablation of *Cdx1*, *Isx*, *Hnf1α*, or *Hnf4α* in mice had no effect on the establishment of the intestinal epithelium (Choi et al., 2006; Garrison et al., 2006; Lee et al., 1998; Pontoglio et al., 1996; Shih et al., 2001; Subramanian et al., 1995). Nevertheless, *Cdx1* (Mutoh et al., 2004), *Isx* (Choi et al., 2006), *HNF1α* (Martin et al., 2000), and *HNF4α* (Garrison et al., 2006) regulate the expression of numerous intestinal genes. Our data support the

In summary, we have identified *Cdx2* as a master regulator in the posterior endoderm, demonstrating that this gene is essential for the establishment of intestinal identity.

EXPERIMENTAL PROCEDURES

Histology, Immunohistochemistry, and Immunofluorescence

Hematoxylin, eosin, and Alcian blue staining was performed in the Morphology Core of the Penn Center for Molecular Studies in Digestive and Liver Diseases. Alkaline phosphatase staining was performed using the Vector Red Alkaline Phosphatase Substrate Kit I (Vector Laboratory, SK-5100). The ABC detection system (Vector Laboratory, PK-6100) was used for immunohistochemistry. Cy2- and Cy3-conjugated fluorescent secondary antibodies were purchased from Jackson Laboratory.

For quantification of intestinal length as well as villi number, length, and width, images of matched control and mutant intestines were analyzed using ImageJ software (NIH). For cell number quantification, BrdU⁺ cells within 50 continuous villi were manually counted from three slides of control and mutant intestines. The percentage of BrdU⁺ cells at each cell position was calculated with the cell located at the bottom of the inter-villus pocket designated as position 0.

Electron Microscopy

Fresh intestinal tissues were washed with PBS and suspended in a fixative solution of 2.5% cacodylate-buffered glutaraldehyde and 4% paraformaldehyde (pH 7.4) for 6 hr. Tissues were rinsed in a cacodylate-buffered solution, postfixed with 2% cacodylate-buffered OsO₄ dehydrated with graded ethanol, clarified in propylene oxide, and embedded in Epon. Seventy-nanometer thin sections were obtained with a Leica UCT ultramicrotome using a Diatome diamond knife and placed on 200 mesh copper grids. Sections were stained with an alcoholic solution of uranyl acetate, followed by a solution of bismuth subnitrate. These sections were examined under a JEOL JEM1010 electron microscope, and digital images were captured using an AMT Advantage HR-Aided Hamamatsu CCD camera. All EM supplies were purchased from Electron Microscopy Sciences, Fort Washington, PA, and Ted Pella, Redding, CA.

Western Blot Analysis

Fresh intestinal tissue lysates were prepared in lysis buffer containing 50 mM Tris (pH 7.5), 150 mM NaCl, 10 mM EDTA, 0.02% Na₃N₃, 50 mM NaF, 1 mM Na₃VO₄, 1% NP40, 1 mM PMSF, and protease inhibitors (Sigma), from E18.5 mouse intestines. 15 µg total lysates were heated at 70°C for 10 min in 4× LDS buffer (Invitrogen) and loaded on 4%–12% SDS-PAGE (Invitrogen). Proteins were transferred to PVDF membranes (Invitrogen). Membranes were stripped in western stripping buffer (Pierce) and reprobed sequentially with corresponding antibodies.

Chromatin Immunoprecipitation

E16.5 control and mutant intestinal tissues were finely minced into small pieces followed by 10 min crosslinking with 1% formaldehyde at 37°C and subjected to chromatin purification and immunoprecipitation as previously described (Rubins et al., 2005) using anti-Cdx2 antibodies (Funakoshi et al., 2008). Input chromatin and ChIP DNA were used as templates in quantitative genomic PCR using an MX3000 PCR machine (Stratagene). The 28S ribosomal genes were used as an internal reference.

Microarray Analysis

One centimeter of the ileum, immediately above the cecum, was dissected from three E18.5 mutant and three control embryos. 250 ng of total RNA was amplified and labeled with Cy3 using the Low Linear Amplification Kit (Agilent Technologies, CA). This labeling reaction produced 1.75–2.0 µg of Cy3-labeled cRNA (antisense), by first converting mRNA primed with an oligo (dT)-T7 primer into dsDNA with MMLV-RT and then amplifying the sample using T7 RNA Polymerase in the presence of Cy3-CTP. After purification, 1.65 µg of cRNA was fragmented and hybridized to the Whole Mouse Genome Oligo Microarray (G4122A; Agilent Technologies, CA) array for 17 hr at 65°C.

Microarray slides were washed and scanned with an Agilent G2565BA Microarray Scanner. Images were analyzed with Feature Extraction 9.5

(Agilent Technologies, CA). Mean foreground intensities were obtained for each spot and imported into the mathematical software package “R.” The Cy3 (green) intensities were corrected for the scanner offset but not further background corrected. The data set was filtered to remove positive control elements. Using the negative controls on the arrays, the background threshold was determined, and all values less than this value were set to the threshold value. Finally, the data were normalized using the Quantile Normalization package in “R” (Bolstad et al., 2003). Complete statistical analysis was then performed in “R” using both the LIMMA and SAM packages to identify statistically significant differential gene expression between the three groups. Microarray data have been deposited in ArrayExpress (www.ebi.ac.uk) under accession number E-MTAB-92.

Quantitative RT-PCR Analysis

Total RNA samples were extracted from E12.5–E18.5 gut tissues. For E12.5 tissues, one biological sample was pooled from two or three guts of like genotype. cDNA synthesis and quantitative RT-PCR analysis were performed as described previously (Gao et al., 2007). qRT-PCR primer sequences are available upon request.

Clustering Analysis and Generation of Heat Map

Hierarchical clustering was performed on the samples (arrays) using the “R” package “pvcust” (Suzuki and Shimodaira, 2006). Additional hierarchical clustering on differentially expressed genes and generation of heat maps were performed using the TM4 Multiple Experiment Viewer software package (Saeed et al., 2003).

Gene Functional Category and Pathway Analysis

Gene functional classification was performed on differentially expressed genes with at least a 4-fold change between control and *Cdx2*-deficient ileum. The Refseq_mRNA IDs of these genes were used for analysis by DAVID Bioinformatics resources, NIH (Dennis et al., 2003). Data were also analyzed through the use of Ingenuity Pathways Analysis (Ingenuity System, www.ingenuity.com) as described previously (Phuc Le et al., 2005).

ACCESSION NUMBERS

Microarray data have been deposited in ArrayExpress (www.ebi.ac.uk) under accession number E-MTAB-92.

SUPPLEMENTAL DATA

Supplemental Data include two tables and eleven figures and can be found with this article online at [http://www.cell.com/developmental-cell/supplemental/S1534-5807\(09\)00084-7](http://www.cell.com/developmental-cell/supplemental/S1534-5807(09)00084-7).

ACKNOWLEDGMENTS

We would like to thank Michael Pack, Joshua Friedman, John Lynch, Blair Madison, Ben Stanger, and Linda Greenbaum for comments on the manuscript; Hong Fu for ES cell culture; Irina Bochkis and Jonathan Schug for help with data analysis and deposition; Karrie Brondell and Elizabeth Helmbrecht for maintaining the mouse colonies; Alan Fox and Olga Smirnova for performing microarray experiments; Kelly Kemnetz for assisting with immunohistochemistry; the Morphology Core of the Penn Center for Molecular Studies in Digestive and Liver Diseases (P30DK50306) for tissue embedding and sectioning; and Neelima Shah and Biomedical Image Core for EM analysis. This work was supported by NIH grants (R01-DK053839 and P01-DK049210) to K.H.K. N.G. is supported by Juvenile Diabetes Research Foundation fellowship 3-2007-521.

Received: August 20, 2008

Revised: December 16, 2008

Accepted: February 18, 2009

Published: April 20, 2009

REFERENCES

- Adaimy, L., Chouery, E., Megarbane, H., Mroueh, S., Delague, V., Nicolas, E., Belguith, H., de Mazancourt, P., and Megarbane, A. (2007). Mutation in WNT10A is associated with an autosomal recessive ectodermal dysplasia: the odontotonycho-dermal dysplasia. *Am. J. Hum. Genet.* **81**, 821–828.
- Adegboyega, P.A., Mifflin, R.C., DiMari, J.F., Saada, J.I., and Powell, D.W. (2002). Immunohistochemical study of myofibroblasts in normal colonic mucosa, hyperplastic polyps, and adenomatous colorectal polyps. *Arch. Pathol. Lab. Med.* **126**, 829–836.
- Aubin, J., Lemieux, M., Tremblay, M., Berard, J., and Jeannotte, L. (1997). Early postnatal lethality in Hoxa-5 mutant mice is attributable to respiratory tract defects. *Dev. Biol.* **192**, 432–445.
- Bates, M.D., Erwin, C.R., Sanford, L.P., Wiginton, D., Bezerra, J.A., Schatzman, L.C., Jegga, A.G., Ley-Ebert, C., Williams, S.S., Steinbrecher, K.A., et al. (2002). Novel genes and functional relationships in the adult mouse gastrointestinal tract identified by microarray analysis. *Gastroenterology* **122**, 1467–1482.
- Beck, F., Erler, T., Russell, A., and James, R. (1995). Expression of Cdx-2 in the mouse embryo and placenta: possible role in patterning of the extra-embryonic membranes. *Dev. Dyn.* **204**, 219–227.
- Beck, F., Chawengsaksophak, K., Waring, P., Playford, R.J., and Furness, J.B. (1999). Reprogramming of intestinal differentiation and intercalary regeneration in Cdx2 mutant mice. *Proc. Natl. Acad. Sci. USA* **96**, 7318–7323.
- Bolstad, B.M., Irizarry, R.A., Astrand, M., and Speed, T.P. (2003). A comparison of normalization methods for high density oligonucleotide array data based on variance and bias. *Bioinformatics* **19**, 185–193.
- Boulet, A.M., and Capecchi, M.R. (1996). Targeted disruption of *hoxc-4* causes esophageal defects and vertebral transformations. *Dev. Biol.* **177**, 232–249.
- Brooke, N.M., Garcia-Fernandez, J., and Holland, P.W. (1998). The *ParaHox* gene cluster is an evolutionary sister of the *Hox* gene cluster. *Nature* **392**, 920–922.
- Charite, J., de Graaff, W., Consten, D., Reijnen, M.J., Korving, J., and Deschamps, J. (1998). Transducing positional information to the *Hox* genes: critical interaction of *cdx* gene products with position-sensitive regulatory elements. *Development* **125**, 4349–4358.
- Chawengsaksophak, K., James, R., Hammond, V.E., Kontgen, F., and Beck, F. (1997). Homeosis and intestinal tumours in Cdx2 mutant mice. *Nature* **386**, 84–87.
- Cheng, P.Y., Lin, C.C., Wu, C.S., Lu, Y.F., Lin, C.Y., Chung, C.C., Chu, C.Y., Huang, C.J., Tsai, C.Y., Korzh, S., et al. (2008). Zebrafish *cdx1b* regulates expression of downstream factors of Nodal signaling during early endoderm formation. *Development* **135**, 941–952.
- Choi, M.Y., Romer, A.I., Hu, M., Lepourcelet, M., Mechoor, A., Yesilaltay, A., Krieger, M., Gray, P.A., and Shivdasani, R.A. (2006). A dynamic expression survey identifies transcription factors relevant in mouse digestive tract development. *Development* **133**, 4119–4129.
- Deb, K., Sivaguru, M., Yong, H.Y., and Roberts, R.M. (2006). Cdx2 gene expression and trophectoderm lineage specification in mouse embryos. *Science* **311**, 992–996.
- Dennis, G., Jr., Sherman, B.T., Hosack, D.A., Yang, J., Gao, W., Lane, H.C., and Lempicki, R.A. (2003). DAVID: Database for Annotation, Visualization, and Integrated Discovery. *Genome Biol.* **4**, P3.
- Deschamps, J., van den Akker, E., Forlani, S., De Graaff, W., Oosterveen, T., Roelen, B., and Roelfsema, J. (1999). Initiation, establishment and maintenance of *Hox* gene expression patterns in the mouse. *Int. J. Dev. Biol.* **43**, 635–650.
- Duprey, P., Chowdhury, K., Dressler, G.R., Balling, R., Simon, D., Guenet, J.L., and Gruss, P. (1988). A mouse gene homologous to the *Drosophila* gene *caudal* is expressed in epithelial cells from the embryonic intestine. *Genes Dev.* **2**, 1647–1654.
- Etensel, B., Temir, G., Karkiner, A., Melek, M., Edirne, Y., Karaca, I., and Mir, E. (2005). Atresia of the colon. *J. Pediatr. Surg.* **40**, 1258–1268.
- Flores, M.V., Hall, C.J., Davidson, A.J., Singh, P.P., Mahagaonkar, A.A., Zon, L.I., Crosier, K.E., and Crosier, P.S. (2008). Intestinal differentiation in zebrafish requires Cdx1b, a functional equivalent of mammalian Cdx2. *Gastroenterology* **135**, 1665–1675.
- Funakoshi, S., Ezaki, T., Kong, J., Guo, R.J., and Lynch, J.P. (2008). Repression of the desmocollin 2 gene expression in human colon cancer cells is relieved by the homeodomain transcription factors Cdx1 and Cdx2. *Mol. Cancer Res.* **6**, 1478–1490.
- Gamer, L.W., and Wright, C.V. (1993). Murine Cdx-4 bears striking similarities to the *Drosophila* *caudal* gene in its homeodomain sequence and early expression pattern. *Mech. Dev.* **43**, 71–81.
- Gao, N., White, P., Doliba, N., Golson, M.L., Matschinsky, F.M., and Kaestner, K.H. (2007). *Foxa2* controls vesicle docking and insulin secretion in mature Beta cells. *Cell Metab.* **6**, 267–279.
- Garrison, W.D., Battle, M.A., Yang, C., Kaestner, K.H., Sladek, F.M., and Duncan, S.A. (2006). Hepatocyte nuclear factor 4alpha is essential for embryonic development of the mouse colon. *Gastroenterology* **130**, 1207–1220.
- Glickman, J.N., Yang, A., Shahsafaei, A., McKeon, F., and Odze, R.D. (2001). Expression of p53-related protein p63 in the gastrointestinal tract and in esophageal metaplastic and neoplastic disorders. *Hum. Pathol.* **32**, 1157–1165.
- Grabin-Botton, A., and Melton, D.A. (2000). Endoderm development: from patterning to organogenesis. *Trends Genet.* **16**, 124–130.
- Hu, Y., Kazenwadel, J., and James, R. (1993). Isolation and characterization of the murine homeobox gene Cdx-1. Regulation of expression in intestinal epithelial cells. *J. Biol. Chem.* **268**, 27214–27225.
- James, R., and Kazenwadel, J. (1991). Homeobox gene expression in the intestinal epithelium of adult mice. *J. Biol. Chem.* **266**, 3246–3251.
- James, R., Erler, T., and Kazenwadel, J. (1994). Structure of the murine homeobox gene *cdx-2*. Expression in embryonic and adult intestinal epithelium. *J. Biol. Chem.* **269**, 15229–15237.
- Kim, B.M., Buchner, G., Miletich, I., Sharpe, P.T., and Shivdasani, R.A. (2005). The stomach mesenchymal transcription factor *Barx1* specifies gastric epithelial identity through inhibition of transient Wnt signaling. *Dev. Cell* **8**, 611–622.
- Kirikoshi, H., Inoue, S., Sekihara, H., and Katoh, M. (2001). Expression of WNT10A in human cancer. *Int. J. Oncol.* **19**, 997–1001.
- Kondo, T., Dolle, P., Zakany, J., and Duboule, D. (1996). Function of posterior *HoxD* genes in the morphogenesis of the anal sphincter. *Development* **122**, 2651–2659.
- Krumlauf, R. (1994). *Hox* genes in vertebrate development. *Cell* **78**, 191–201.
- Lau, S.T., and Caty, M.G. (2006). Hindgut abnormalities. *Surg. Clin. North Am.* **86**, 301–316.
- Lee, Y.H., Sauer, B., and Gonzalez, F.J. (1998). Laron dwarfism and non-insulin-dependent diabetes mellitus in the *Hnf-1alpha* knockout mouse. *Mol. Cell. Biol.* **18**, 3059–3068.
- Lee, C.S., Friedman, J.R., Fulmer, J.T., and Kaestner, K.H. (2005). The initiation of liver development is dependent on *Foxa* transcription factors. *Nature* **435**, 944–947.
- Li, X., Madison, B.B., Zacharias, W., Kolterud, A., States, D., and Gumucio, D.L. (2007). Deconvoluting the intestine: molecular evidence for a major role of the mesenchyme in the modulation of signaling cross talk. *Physiol. Genomics* **29**, 290–301.
- Macdonald, P.M., and Struhl, G. (1986). A molecular gradient in early *Drosophila* embryos and its role in specifying the body pattern. *Nature* **324**, 537–545.
- Madison, B.B., Braunstein, K., Kuizon, E., Portman, K., Qiao, X.T., and Gumucio, D.L. (2005). Epithelial hedgehog signals pattern the intestinal crypt-villus axis. *Development* **132**, 279–289.
- Manley, N.R., and Capecchi, M.R. (1995). The role of *Hoxa-3* in mouse thymus and thyroid development. *Development* **121**, 1989–2003.
- Martin, M.G., Wang, J., Solorzano-Vargas, R.S., Lam, J.T., Turk, E., and Wright, E.M. (2000). Regulation of the human Na(+)-glucose cotransporter

- gene, SGLT1, by HNF-1 and Sp1. *Am. J. Physiol. Gastrointest. Liver Physiol.* 278, G591–G603.
- McGinnis, W., and Krumlauf, R. (1992). Homeobox genes and axial patterning. *Cell* 68, 283–302.
- Meyer, B.I., and Gruss, P. (1993). Mouse Cdx-1 expression during gastrulation. *Development* 117, 191–203.
- Mlodzik, M., Fjose, A., and Gehring, W.J. (1985). Isolation of caudal, a Drosophila homeo box-containing gene with maternal expression, whose transcripts form a concentration gradient at the pre-blastoderm stage. *EMBO J.* 4, 2961–2969.
- Moreno, E., and Morata, G. (1999). Caudal is the Hox gene that specifies the most posterior Drosophila segment. *Nature* 400, 873–877.
- Mutoh, H., Sakurai, S., Satoh, K., Osawa, H., Hakamata, Y., Takeuchi, T., and Sugano, K. (2004). Cdx1 induced intestinal metaplasia in the transgenic mouse stomach: comparative study with Cdx2 transgenic mice. *Gut* 53, 1416–1423.
- Ouban, A., Nawab, R.A., and Coppola, D. (2002). Diagnostic and pathogenetic implications of colorectal carcinomas with multidirectional differentiation: a report of 4 cases. *Clin. Colorectal Cancer* 1, 243–248.
- Phuc Le, P., Friedman, J.R., Schug, J., Brestelli, J.E., Parker, J.B., Bochkis, I.M., and Kaestner, K.H. (2005). Glucocorticoid receptor-dependent gene regulatory networks. *PLoS Genet.* 1, e16.
- Pitera, J.E., Smith, V.V., Thorogood, P., and Milla, P.J. (1999). Coordinated expression of 3' hox genes during murine embryonal gut development: an enteric Hox code. *Gastroenterology* 117, 1339–1351.
- Pollock, R.A., Jay, G., and Biebrich, C.J. (1992). Altering the boundaries of Hox3.1 expression: evidence for antipodal gene regulation. *Cell* 71, 911–923.
- Pontoglio, M., Barra, J., Hadchouel, M., Doyen, A., Kress, C., Bach, J.P., Babinet, C., and Yaniv, M. (1996). Hepatocyte nuclear factor 1 inactivation results in hepatic dysfunction, phenylketonuria, and renal Fanconi syndrome. *Cell* 84, 575–585.
- Que, J., Okubo, T., Goldenring, J.R., Nam, K.T., Kurotani, R., Morrissey, E.E., Taranova, O., Pevny, L.H., and Hogan, B.L. (2007). Multiple dose-dependent roles for Sox2 in the patterning and differentiation of anterior foregut endoderm. *Development* 134, 2521–2531.
- Roberts, D.J. (2000). Molecular mechanisms of development of the gastrointestinal tract. *Dev. Dyn.* 219, 109–120.
- Rodriguez, C.I., Buchholz, F., Galloway, J., Sequerra, R., Kasper, J., Ayala, R., Stewart, A.F., and Dymecki, S.M. (2000). High-efficiency deleter mice show that FLP is an alternative to Cre-loxP. *Nat. Genet.* 25, 139–140.
- Rubins, N.E., Friedman, J.R., Le, P.P., Zhang, L., Brestelli, J., and Kaestner, K.H. (2005). Transcriptional networks in the liver: hepatocyte nuclear factor 6 function is largely independent of Foxa2. *Mol. Cell. Biol.* 25, 7069–7077.
- Saeed, A.I., Sharov, V., White, J., Li, J., Liang, W., Bhagabati, N., Braisted, J., Klapa, M., Currier, T., Thiagarajan, M., et al. (2003). TM4: a free, open-source system for microarray data management and analysis. *Biotechniques* 34, 374–378.
- Schroder, N., Sekhar, A., Geffers, I., Muller, J., Dittrich-Breiholz, O., Kracht, M., Wedemeyer, J., and Gossler, A. (2006). Identification of mouse genes with highly specific expression patterns in differentiated intestinal epithelium. *Gastroenterology* 130, 902–907.
- Shih, D.Q., Bussen, M., Sehayek, E., Ananthanarayanan, M., Shneider, B.L., Suchy, F.J., Shefer, S., Bollilini, J.S., Gonzalez, F.J., Breslow, J.L., and Stoffel, M. (2001). Hepatocyte nuclear factor-1alpha is an essential regulator of bile acid and plasma cholesterol metabolism. *Nat. Genet.* 27, 375–382.
- Shimizu, T., Bae, Y.K., and Hibi, M. (2006). Cdx-Hox code controls competence for responding to Fgfs and retinoic acid in zebrafish neural tissue. *Development* 133, 4709–4719.
- Silberg, D.G., Swain, G.P., Suh, E.R., and Traber, P.G. (2000). Cdx1 and cdx2 expression during intestinal development. *Gastroenterology* 119, 961–971.
- Silberg, D.G., Sullivan, J., Kang, E., Swain, G.P., Moffett, J., Sund, N.J., Sackett, S.D., and Kaestner, K.H. (2002). Cdx2 ectopic expression induces gastric intestinal metaplasia in transgenic mice. *Gastroenterology* 122, 689–696.
- Skromne, I., Thorsen, D., Hale, M., Prince, V.E., and Ho, R.K. (2007). Repression of the hindbrain developmental program by Cdx factors is required for the specification of the vertebrate spinal cord. *Development* 134, 2147–2158.
- Stoffers, D.A., Zinkin, N.T., Stanojevic, V., Clarke, W.L., and Habener, J.F. (1997). Pancreatic agenesis attributable to a single nucleotide deletion in the human IPF1 gene coding sequence. *Nat. Genet.* 15, 106–110.
- Subramanian, V., Meyer, B.I., and Gruss, P. (1995). Disruption of the murine homeobox gene Cdx1 affects axial skeletal identities by altering the mesodermal expression domains of Hox genes. *Cell* 83, 641–653.
- Suzuki, R., and Shimodaira, H. (2006). Pvcust: an R package for assessing the uncertainty in hierarchical clustering. *Bioinformatics* 22, 1540–1542.
- Tamai, Y., Nakajima, R., Ishikawa, T., Takaku, K., Seldin, M.F., and Taketo, M.M. (1999). Colonic hamartoma development by anomalous duplication in Cdx2 knockout mice. *Cancer Res.* 59, 2965–2970.
- van den Akker, E., Forlani, S., Chawengsaksophak, K., de Graaff, W., Beck, F., Meyer, B.I., and Deschamps, J. (2002). Cdx1 and Cdx2 have overlapping functions in anteroposterior patterning and posterior axis elongation. *Development* 129, 2181–2193.
- van Nes, J., de Graaff, W., Lebrin, F., Gerhard, M., Beck, F., and Deschamps, J. (2006). The Cdx4 mutation affects axial development and reveals an essential role of Cdx genes in the ontogenesis of the placental labyrinth in mice. *Development* 133, 419–428.
- Wang, Y., Yabuuchi, A., McKinney-Freeman, S., Ducharme, D.M., Ray, M.K., Chawengsaksophak, K., Archer, T.K., and Daley, G.Q. (2008). Cdx gene deficiency compromises embryonic hematopoiesis in the mouse. *Proc. Natl. Acad. Sci. USA* 105, 7756–7761.
- Warot, X., Fromental-Ramain, C., Fraulob, V., Chambon, P., and Dolle, P. (1997). Gene dosage-dependent effects of the Hoxa-13 and Hoxd-13 mutations on morphogenesis of the terminal parts of the digestive and urogenital tracts. *Development* 124, 4781–4791.
- Wells, J.M., and Melton, D.A. (1999). Vertebrate endoderm development. *Annu. Rev. Cell Dev. Biol.* 15, 393–410.
- Wolgemuth, D.J., Behringer, R.R., Mostoller, M.P., Brinster, R.L., and Palmiter, R.D. (1989). Transgenic mice overexpressing the mouse homeobox-containing gene Hox-1.4 exhibit abnormal gut development. *Nature* 337, 464–467.
- Yang, Q., Bermingham, N.A., Finegold, M.J., and Zoghbi, H.Y. (2001). Requirement of Math1 for secretory cell lineage commitment in the mouse intestine. *Science* 294, 2155–2158.
- Zacchetti, G., Duboule, D., and Zakany, J. (2007). Hox gene function in vertebrate gut morphogenesis: the case of the caecum. *Development* 134, 3967–3973.
- Zakany, J., and Duboule, D. (1999). Hox genes and the making of sphincters. *Nature* 401, 761–762.